


GRB Discoveries with Swift

Neil Gehrels
NASA-GSFC


Swift GRB 070420

BAT prompt emission



3 instruments, each with:
- lightcurves
- images
- spectra

XRT afterglow



Long GRBs

**63 Swift Long
GRB Redshifts**


6.29	050904	2.35	070110
6.47	060927	2.31	070506
6.3	050814	2.30	060124
5.11	060522	2.20	050922C
4.9	060510B	2.17	070810
4.41	060223A	2.04	070611
4.27	050505	1.95	050315
4.05	060206	1.71	050802
3.97	050730	1.55	051111
3.91	060210	1.51	060502A
3.71	060605	1.50	070306
3.69	060906	1.49	060418
3.62	070721B	1.44	050318
3.53	060115	1.31	061121
3.44	061105	1.29	050126
3.43	060707	1.26	061007
3.36	061222B	1.17	070208
3.34	050908	0.97	070419A
3.24	050319	0.94	051014B
3.21	060926	0.84	070118
3.21	060526	0.83	050824
3.08	060607A	0.76	06110A
2.95	070411	0.70	060904B
3.90	050401	0.65	050416A
2.82	050603	0.62	070612A
2.71	060714	0.61	050525A
2.58	060604	0.54	060729
2.61	050820A	0.44	060512
2.50	070529	0.125	060614
2.45	070802	0.089	060505
2.43	060908	0.033	060218
2.35	051109A		

z	GRB	Optical/IR Brightness
6.29	050904	J = 18 @ 3 hrs
5.6	060927	I = 16 @ 2 min
5.3	050814	K = 18 @ 23 hrs
5.11	060522	R = 21 @ 1.5 hrs

GRB Host Spectroscopy


GRB 050505
 $z = 4.275$
 Damped Ly α
 $N(\text{H}) = 10^{22} \text{ cm}^{-2}$
 $n \sim 10^2 \text{ cm}^{-3}$
 $Z = 0.06 Z_{\odot}$
 $M_{\text{progenitor}} < 25 M_{\odot}$

Metallicity vs Redshift




Savaglio 2006

GRB 050904 $z=6.29$






Subaru Telescope
 Kawai et al. 2006

GRB 060218: GRB + Supernova




Super-long GRB - ~35 minutes
 BAT, XRT, UVOT during GRB
 $z = 0.033$ $d = 145 \text{ Mpc}$
 SN 2006aj SN Ib/c
 $E_{100} = \text{few} \times 10^{49} \text{ erg}$ - underluminous

Afterglows



Short GRBs

Short GRB Time Structure



Short GRB - Current Status

Swift short GRB observations

- 23 short bursts detected (+ 2 from HETE, +1 from INTEGRAL)
- 78% with X-ray afterglow detected by XRT (95% long GRBs)
- 28% with optical detection (58% long GRBs)
- ~50% with host IDs

~1/2 shorts accompanied by soft extended emission up to 100 sec


Redshift range from $z = 0.2$ to 1

- $\langle z \rangle_{\text{short}} = 0.6$
- $\langle z \rangle_{\text{long}} = 2.3$

GRB 070714B $z = 0.92$
(Graham et al. 2007)

3 Types of GRBs

Swift GRBs (mostly)



Implications for Grav. Wave Detections

Assuming all short GRBs are due to NS-NS mergers, merger rate is $\sim 300 \text{ Gpc}^{-3} \text{ yr}^{-1}$

[Consistent with NS-NS population synthesis modeling O'Shaughnessy, Kalogera, & Belczynski (2005)]

\Rightarrow Advanced LIGO detection rate of $\sim 30 \text{ yr}^{-1}$

Nakar et al.:

Possible much higher rates of $10^5 \text{ Gpc}^{-3} \text{ yr}^{-1}$.

\Rightarrow Detection with enhance LIGO

Swift will be in orbit until > 2020

